


My Profile


[Home](#) / [Chemistry](#) / [Pharmaceutical and Medicinal Chemistry](#)

▶ [HOME](#)
 ▶ [ABOUT US](#)
 ▶ [CONTACT US](#)
 ▶ [HELP](#)

**Journal of Pharmaceutical Sciences****Volume 73, Issue 4, Pages 468 - 471**

Published Online: 18 Sep 2006

Copyright © 1984 Wiley-Liss, Inc., A Wiley Company

☐ [Save Title to My Profile](#)
☐ [Set E-Mail Alert](#)


Go to the homepage for this journal to access trials, sample copies, editorial and author information, news, and more. ▶

☐ e-mail ☐ print

Published on behalf of

[Go to Society Site](#)SEARCH ☒ All Content☐ Publication[Advanced Search](#)[CrossRef / Google Search](#)[Acronym Finder](#)

SEARCH IN THIS TITLE

Journal of Pharmaceutical Sciences

☒ [Save Article to My Profile](#)
☒ [Download Citation](#)
[< Previous Abstract](#) | [Next Abstract >](#)
[Abstract](#) | [References](#) | [Full Text: PDF \(480k\)](#) | [Related Articles](#) | [Citation Tracking](#)

All Fields

Research Article**Viscosity of xanthan gum solutions at low shear rates**

Joel L. Zatz*, Steven Knapp

Rutgers College of Pharmacy, Piscataway, NJ 08854

*Correspondence to Joel L. Zatz, Rutgers College of Pharmacy, Piscataway, NJ 08854

Funded by:

• KELTROL® xanthan gum, and Haake, Inc.

KEYWORDS

Viscosity - xanthan gum in aqueous solutions, low shear rates, effect of added salts • Xanthan gum - aqueous solutions, viscosity at low shear rates, effect of added salts • Shear rate - low range, viscosity of xanthan gum in aqueous solutions, effect of added salts

ABSTRACT

The viscosity of xanthan gum solutions in the low shear region was investigated with the aid of a Couette instrument. All solutions were highly pseudoplastic. Solutions containing 0.3-0.5% of the gum exhibited a highly ordered phase at very low shear. Viscosity, the degree of pseudoplasticity, and the value of the transition from soft gel to pseudoplastic behavior were directly related to gum concentration. The effect of the addition of a salt on viscosity depended on the xanthan gum concentration. The viscosity of a 0.3% xanthan gum solution was practically unaffected by the salts. Higher gum concentrations exhibited a viscosity increase when salt was present. Concentrations >0.3% exhibited a viscosity decrease in the presence of a salt. All viscosity effects seemed to reach limiting values at $\sim 10^{-3}$ to 3.3×10^{-3} M salt. No major differences were observed between sodium chloride, calcium chloride, and sodium citrate in their influence on xanthan gum viscosity.

Received: 3 September 1982; Accepted: 30 November 1982

DIGITAL OBJECT IDENTIFIER (DOI)10.1002/jps.2600730410 [About DOI](#)

SEARCH BY CITATION

Vol: Issue: Page:

INDEXED BY



This journal is listed in the National Library of Medicine PubMed index.

SIGN UP NOW

**MOBILE EDITION**

Get select content from Wiley's leading publications delivered to your PDA —

[Sign up now](#)

REPRINT INQUIRIES

